

References

1. Hayes EB. Zika virus outside Africa. *Emerg Infect Dis.* 2009;15:1347–50. <http://dx.doi.org/10.3201/eid1509.090442>

2. Lanciotti RS, Kosoy OL, Laven JJ, Velez JO, Lambert AJ, Johnson AJ, et al. Genetic and serologic properties of Zika virus associated with an epidemic, Yap State, Micronesia, 2007. *Emerg Infect Dis.* 2008;14:1232–9. <http://dx.doi.org/10.3201/eid1408.080287>

3. European Centre for Disease Prevention and Control. Zika virus—Pacific (08). *ProMed.* 2014 Mar 16. <http://www.promedmail.org>, archive no. 2014 0316.2335754.

4. Schwan K. Zika virus—Pacific (07). *ProMed.* 2014 Mar 9. <http://www.promedmail.org>, archive no. 20140309.2322907.

5. Tappe D, Schmidt-Chanasit J, Ries A, Ziegler U, Müller A, Stich A. Ross River virus infection in a traveller returning from northern Australia. *Med Microbiol Immunol (Berl).* 2009;198:271–3. <http://dx.doi.org/10.1007/s00430-009-0122-9>

6. Tappe D, Rissland J, Gabriel M, Emmerich P, Gunther S, Held G, et al. First case of laboratory-confirmed Zika virus infection imported to Europe, November 2013. *Euro Surveill.* 2014;19:20685.

7. Fonseca K. Zika virus—Canada ex Thailand. *ProMed.* 2013 May 29. <http://www.promedmail.org>, archive no. 2013 0529.1744108.

8. Kwong JC, Druce JD, Leder K. Zika virus infection acquired during brief travel to Indonesia. *Am J Trop Med Hyg.* 2013;89:516–7. <http://dx.doi.org/10.4269/ajtmh.13-0029>

9. Foy BD, Kobylinski KC, Chilson Foy JL, Blitvich BJ, Travassos da Rosa A, Haddow AD, et al. Probable non-vector-borne transmission of Zika virus, Colorado, USA. *Emerg Infect Dis.* 2011;17:880–2. <http://dx.doi.org/10.3201/eid1705.101939>

10. Kutsuna S, Kato Y, Takasaki T, Moi ML, Kotaki A, Uemura H. Two cases of Zika virus imported from French Polynesia to Japan, December 2013 to January 2014. *Euro Surveill.* 2014;19:pi=20683.

Address for correspondence: Jonas Schmidt-Chanasit, Bernhard Nocht Institute for Tropical Medicine, WHO Collaborating Centre for Arbovirus and Haemorrhagic Fever Reference and Research, Bernhard-Nocht-Strasse 74, 20359 Hamburg, Germany; email: jonassi@gmx.de

SEARCH PAST ISSUES OF EID AT
WWW.CDC.GOV/EID

***Yersinia pestis* in
Pulex irritans
Fleas during
Plague Outbreak,
Madagascar**

To the Editor: *Yersinia pestis* (family *Enterobacteriaceae*) is a bacterium that can cause high rates of death in susceptible mammals and can provoke septicemic, pneumonic, and bubonic plague in humans (1). This zoonotic pathogen can be transmitted directly by infectious droplets or by contact with contaminated fluid or tissue or indirectly through flea bites (1). Plague was introduced into Madagascar in 1898 from rat-infested steamships that had sailed from affected areas (2). Now, Madagascar is 1 of 2 countries in Africa that have reported cases of human plague every year since 1991 (3). During January 2008–January 2013, the number of human plague cases reported in Madagascar ranged from 312 to 648 per year. Of these, 61.8%–75.5% were laboratory confirmed (National Plague Laboratory of the Ministry of Health, pers. comm.). Most (>83%) confirmed cases were bubonic plague, which most commonly results from flea bites, suggesting that these bites were the most common mode of *Y. pestis* transmission. In Madagascar, *Xenopsylla cheopis* fleas have been known as the primary plague vector in urban areas, whereas *Synopsyllus fonquerniei* fleas have been usually involved in plague transmission in rural areas (2).

In January 2013, a total of 9 suspected bubonic plague cases, 3 confirmed, were reported in Soavina, a

rural area in the district of Ambatofinandra, Madagascar. Domestic fleas were collected with candle traps inside 5 houses during 3 nights (Table). Fleas were also caught on small mammals trapped inside houses and outside in the sisal fences and rice fields (Table). A total of 319 fleas belonging to 5 species in 5 genera were collected inside and outside the houses, an average of 44 per house (maximum 71): *Pulex irritans*, *Echidnophaga gallinacea*, and *Ctenocephalides canis* fleas were collected inside the houses (244, 76.5%), and *S. fonquerniei* and *X. cheopis* fleas were collected outside (75, 23.5%). The human flea, *P. irritans*, was the most collected flea species (233, 73.3%), followed by *S. fonquerniei* (62, 19.4%), *X. cheopis* (13, 4.1%), *E. gallinacea* (10, 3.1%), and *C. canis* (1, 0.3%).

Bacterial DNA was extracted from 277 fleas of 5 species: 233 *P. irritans*, 24 *S. fonquerniei*, 9 *X. cheopis*, 10 *E. gallinacea*, and 1 *C. canis*. PCR to detect *Y. pestis* was performed by using primers YP1 (5'-ATC TTA CTT TCC GTG AGA AG-3') and YP2 (5'-CTT GGA TGT TGA GCT TCC TA-3') to amplify a 478-bp fragment (4). *Y. pestis* DNA was then amplified and genotyped by Beckman Coulter Genomics Inc. (Takeley, United Kingdom). The positive control was *Y. pestis* reference strain (strain 6/69, 3 × 10⁸ bacteria/mL; Institut Pasteur de Madagascar).

Detection of *Y. pestis* was carried out on 274 fleas belonging to 5 flea species: 230 *P. irritans* (181 unfed and 49 engorged), 24 *S. fonquerniei* (15 unfed and 9 engorged), 9 *X. cheopis* (8 unfed and 1 engorged), 10 *E. gallinacea* (blood-feeding status not identified), and 1 unfed *C. canis*. *Y. pestis*

Table. Fleas collected inside and outside houses in Soavina, Madagascar, January 2013			
Species	Total no. (%)	No. (%) inside	No. (%) outside
<i>Pulex irritans</i>	233 (73.0)	233 (95.5)	0
<i>Ctenocephalides canis</i>	1 (0.3)	1 (0.4)	0
<i>Echidnophaga gallinacea</i>	10 (3.1)	10 (4.1)	0
<i>Synopsyllus fonquerniei</i>	62 (19.4)	0	62 (82.7)
<i>Xenopsylla cheopis</i>	13 (4.1)	0	13 (17.3)
Total	319 (100)	244 (100)	75 (100)

was detected in 9 *P. irritans* fleas (7 male [6 unfed and 1 engorged] and 2 [engorged] female) from 3 houses, including the house where a confirmed human case of plague had occurred (online Technical Appendix, <http://wwwnc.cdc.gov/EID/article/20/8/13-0629-Techapp1.pdf>). Eight sequences (GenBank accession nos. KJ361938–KJ361945) were obtained and share 99% nucleotide homology with plasminogen activator genes of *Y. pestis* published in GenBank (accession nos. AF528537, AY305870). No *Y. pestis* was detected in the 24 *S. fonquerniei*, 9 *X. cheopis*, 10 *E. gallinacea*, or 1 *C. canis* fleas collected.

Although only *X. cheopis* and *S. fonquerniei* fleas had previously been described as plague vectors in Madagascar, *P. irritans* fleas were most commonly collected during this field study; engorged and unfed male and female *P. irritans* fleas carried *Y. pestis*. Other studies have found *P. irritans* fleas in the plague risk area in other countries in Africa (5,6); one study found that *P. irritans* fleas may play a role in plague epidemiology in Tanzania (5). Data on *P. irritans* fleas in rats make it unlikely that these fleas are involved in rat-to-human transmission of *Y. pestis* in Madagascar. During 1922–1995, a total of 118,608 rats were caught and examined in Madagascar, but only 148 *P. irritans* fleas were identified, and none have been found on rats since 1996 (<http://www.pasteur.mg/spip.php?rubrique124>). The high density of *P. irritans* fleas we observed in villages where plague outbreaks occurred in late 2012 and early 2013 (<http://www.pasteur.mg/spip.php?rubrique124>) supports the possibility that *P. irritans* fleas played a role in domestic human-to-human transmission of *Y. pestis* during these outbreaks.

**Jocelyn Ratovonjato,
Minoarisoa Rajerison,
Soanandrasana Rahelinirina,
and Sébastien Boyer**


Author affiliations: Institut Pasteur de Madagascar, Antananarivo, Madagascar (J. Ratovonjato, S. Boyer); and World Health Organization Collaborating Centre, Institut Pasteur de Madagascar, Antananarivo (M. Rajerison, S. Rahelinirina)

DOI: <http://dx.doi.org/10.3201/eid2008.130629>

References

1. Bulter T. Plague and other *Yersinia* infections. In: Greenough WB III, Merigan TC, editors. Current topics in infectious disease. New York: Plenum; 1983. p. 71–92.
2. Brygoo ER. Epidemiology of the plague at Madagascar [in French]. Arch Inst Pasteur Madagascar. 1966;35:7–147.
3. World Health Organization. Report on global surveillance of epidemic-prone infectious diseases—plague [cited 2013 Apr 15]. http://www.who.int/csr/resources/publications/plague/CSR_ISR_2000_1/en/index5.html
4. Hinnebusch J, Schwan TG. New method for plague surveillance using polymerase chain reaction to detect *Yersinia pestis* in fleas. J Clin Microbiol. 1993;31:1511–4.
5. Laudisoit A, Leirs H, Makundi RH, Van Dongen S, Davis S, Neerincx S, et al. Plague and the human flea, Tanzania. Emerg Infect Dis. 2007;13:687–93. <http://dx.doi.org/10.3201/eid1305.061084>
6. Eisen RJ, Gage KL. Transmission of flea-borne zoonotic agents. Annu Rev Entomol. 2012;57:61–82. <http://dx.doi.org/10.1146/annurev-ento-120710-100717>

Address for correspondence: Jocelyn Ratovonjato, Medical Entomology Unit, Institut Pasteur de Madagascar, BP 1274, Antananarivo 101, Madagascar; email: ratov@pasteur.mg



Serologic Surveillance for West Nile Virus in Dogs, Africa

To the Editor: West Nile fever is caused by the West Nile virus (WNV), a mosquito-borne member of the genus *Flavivirus*. Birds are the natural reservoir of the virus, which is maintained in nature in a mosquito–bird–mosquito transmission cycle. WNV has been detected in many regions worldwide, including North America, Europe, Africa, the Near East, and Asia (1). WNV has been shown to cause meningoencephalitis in humans and horses. In the United States, seroconversion in dogs was detected 6 weeks before a human case was reported (2). Thus, dogs could be considered as sentinels for WNV infection, but their role as reservoir is unlikely because of short-term and low levels of viremia (3). In this study, we determined the seroprevalence of WNV in dogs living close to humans in different environments to assess their role as sentinels of this potentially severe zoonosis.

During 2003–2012, blood samples were collected from 753 adult dogs from France and 6 countries in Africa (Table). Samples were centrifuged within 24 h after collection, separated, frozen at -20°C , and sent to the virology laboratory of the Institut de Recherche Biomédicale des Armées (Marseille, France). Each sample was systematically tested for IgG against WNV by using an in-house ELISA with inactivated WNV as antigen. Serum samples were considered positive if the optical density at 450 nm was >3 -fold the mean of that for negative antigen. Because of the antigenic cross-reactivity among flaviviruses, all positive samples were further tested by Western blot for WNV-specific antibodies (4); seroprevalence was calculated on the basis of Western blot–confirmed cases only.